

MAR 24 2005

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 23.Mar.05		3. REPORT TYPE AND DATES COVERED MAJOR REPORT
4. TITLE AND SUBTITLE RECOMBINANT PHAGE PROBES FOR SALMONELLA TYPHIMURIUM DETECTION			5. FUNDING NUMBERS	
6. AUTHOR(S) MAJ OLSEN ERIC V				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) AUBURN UNIVERSITY MAIN CAMPUS			8. PERFORMING ORGANIZATION REPORT NUMBER CI04-1010	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) THE DEPARTMENT OF THE AIR FORCE AFIT/CIA, BLDG 125 2950 P STREET WPAFB OH 45433			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION AVAILABILITY STATEMENT Unlimited distribution In Accordance With AFI 35-205/AFIT Sup 1			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words)				
14. SUBJECT TERMS			15. NUMBER OF PAGES 3	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT	

Name of submitting author: Eric Olsen.

Degrees: A.S., *Galveston College, Galveston, TX*; B.S. Medical Technology, *University of Texas, Galveston, TX*; M.S. Technical and Occupational Education, *University of Southern Mississippi, Long Beach, MS*; M.S. Microbiology, *Auburn University, Auburn, AL*.

Certifications: MT (ASCP), CLS (NCA)

Current positions: Ph.D. student, Department of Biological Sciences, Auburn University; active duty US Air Force Laboratory Officer (Major) with specialties in laboratory medicine (43T3A) and microbiology (43T3B).

Institutions: Auburn University, Auburn, AL; Air Force Institute of Technology, Wright-Patterson AFB, OH.

✓ Mailing address: 435 Bama Park Road, Dadeville, AL, USA, 36853.

✓ Phone: 334-319-3804 (CP).

E-mail: olsenev@auburn.edu

Disclaimer: The views expressed in this article are those of the author and do not reflect the official policy or position of the United States Air Force, Department of Defense, or the U.S. Government.

Type of abstract: Oral, Poster (* = **presenting author**).

RECOMBINANT PHAGE PROBES FOR *SALMONELLA TYPHIMURIUM* DETECTION

Eric Olsen, M.S.^{1*}, Iryna Sorokulova, Ph.D.², I - Hsuan Chen, M.S.², James Barbaree, Ph.D.¹, Vitaly Vodyanoy, Ph.D.³, and Valery A. Petrenko, Ph.D.²

¹Auburn University, Department of Biological Sciences, Auburn AL 36849. ²Auburn University, College of Veterinary Medicine, Department of Pathobiology, Auburn AL 36849. ³Auburn University, College of Veterinary Medicine, Department of Anatomy, Physiology and Pharmacology, Auburn AL 36849.

Salmonella typhimurium is a leading cause of inadvertent gastrointestinal foodborne illness in the United States. Although few actual accounts of deliberate food contamination have been documented in the United States, the recent advent of biocrimes and terrorism in our country suggests that this trend will not continue, highlighting the importance of rapidly identifying biological agents, regardless of the contamination origin, as one part of a comprehensive strategic plan to secure the public food supply. There is an urgent need for deployable, real-time threat agent detectors to replace traditional methods of food safety analysis that are slower, labor-intensive, and cost-inefficient. Confirmation of presence in food products can take as long as 48 hours by conventional culture. Current rapid detection initiatives include biosensors that routinely incorporate antibodies as the biorecognition unit. Although sensitive and specific, antibodies are costly and may degrade under unfavorable environmental conditions. We believe that a stable, inexpensive substitute for antibodies is filamentous phage manipulated through phage display technique then affinity selected for specificity to *S. typhimurium* from billion-clone phage landscape libraries. Our results show that recombinant phage affinity selected against *S. typhimurium* can be 12,000-22,000 times for more specific than controls and 10-1000 times more selective for *S. typhimurium* than

20050328 018

other select enterobacteria. We anticipate that these highly specific, selective phage binders will build upon our current biosensor development initiatives for the rapid detection of biological agents such as *S. typhimurium*.


Binding Confirmation

Methods


Phage specificity

Phage selectivity




Phage selectivity



Selection of *S. typhimurium* specific phage clones from the phage library was performed by affinity selection in solution. Following heat incubation (70° C) and depletion of aggregates by precipitation at high centrifugal speed, a phage library was added to a suspension of *S. typhimurium* for 1 hour to effect cell specific binding. Cells were washed 10X to remove any non-specific binders. Cell bound phage was collected from both eluate and lysate of cells, then amplified in *E. coli*. DNA of recovered phage clones was used to determine the amino acid sequence of their capsid proteins. PvuII digested templates



Selection of *S. typhimurium* specific phage clones from the phage library was performed by affinity selection in solution. Following heat incubation (70° C) and depletion of aggregates by centrifugation at high centrifugal speed, a phage library was added to a suspension of *S. typhimurium* for 1 hour to effect cell specific binding. Cells were washed 10X to remove any non-specific binding. Cell bound phage was collected from both eluate and wash fractions. Cells were lysed with 100 µl of 1% Sarkosyl NL30. Lysate of cells, then amplified in *E. coli*. DNA of recovered phage clones was used to determine the amino acid sequence of their respective PVIIL displayed peptides.



Figs. 4-6. Dose-dependent binding of E2 to *S. typhimurium*. Determination of phage specificity and selectivity by precipitation assay, initially mediated at a concentration of 10^6 CFU/ml, resulted in low absolute yield percentage (= 2% for *S. typhimurium* prior to yield normalization). To determine if dose-dependent concentration was a factor affecting binding, titration of E2 in conjunction with precipitation assay was performed by substituting the single concentration of E2 utilized with concentrations ranging from 10^{12} to 10^1 CFU/ml. A higher yield was demonstrated at a concentration of E2 utilized with concentrations ranging from 10^{12} to 10^4 CFU/ml, when ratios of phage to cells approached 1.0, suggesting a "stoichiometric" mode of phage-bacteria interaction. Binding isotherm (Figure 5) confirms concentration-dependent binding of phage to cells in solution. The K_d apparent of the complex calculated from the Hill plot is 3.18×10^9 CFU/ml (= 0.01 nanomolar). Hill coefficient ($n = 0.91$) estimated from Hill plot (Figure 6) is in good agreement with yield results.



Fig. 3. FACS analysis of phage E2 binding to *S. typhimurium*. Fluorescence associated with cells treated with phage is greater than that of untreated cells and is dependent upon the concentration of phage utilized.




Fig. 2. Precipitation assay demonstrated 90% greater affinity of phage E2 (mean yield normalized) for *S. typhimurium* in comparison to challenge bacteria. Mean yield % is an average of 3 separate experiments normalized to a maximal mean yield of 2.8% from *S. typhimurium*.

typhimurium

Results




Fig. 1. Specificity of select phage (10^6 CFU/ml) were confirmed for *S. typhimurium* by phage capture, in comparison to ELISA (data not shown), which uses a solid support. Binding of best clones to *S. typhimurium* were 12,000–22,000 times greater than wild-type control phage 18-5, with phage clone E2 (VTPPTQHQ) possessing the highest binding efficiency among all select phage tested.

Conclusions

These results confirm our group's previous research efforts in the development of phage probes for the detection of biological molecules, commending landscape phage as substitute antibodies. We anticipate that these highly specific, selective phage binders will build upon our current biosensor development initiatives for the rapid detection of biological agents such as *S. typhimurium*.

Acknowledgements

This work was supported by USDA grant No. 20013439410295C, and ARO/DARPA grant # DAAD 19-01-10454